

IROQUOIS CENTRAL SCHOOL DISTRICT

HS FAX: 716-995-2449, MS FAX: 716-995-2455, ELMA FAX: 716-805-7040, MARILLA FAX: 716-805-7046, WALES FAX: 716-805-7075
Phone: 716- 652-3000, HS ext. 7600, MS ext. 6600, Elma ext. 2600, Marilla ext. 3600, Wales ext. 4600

LIFE THREATENING ALLERGY – EMERGENCY CARE PLAN

20 – 20 School Year

Student Name: _____ DOB _____ Grade _____

Identified Allergen: _____

Asthma Yes No Other relevant health concern: _____

<i>Contact Information</i>
Mother: _____
Father: _____
Emergency Contact: _____
Additional Contact: _____

School Nurse _____ Phone extension _____

If nurse cannot be reached, call: Main Office at extension _____ or 911

AN ALLERGIC REACTION MAY INCLUDE ANY OR ALL OF THESE SYMPTOMS

- ⊛ **General:** Dizziness, loss of consciousness, feeling of panic or doom
- ⊛ **Mouth:** Swelling of lips, face, tongue, throat; a report that the mouth "feels hot"
- ⊛ **Breathing:** Wheezing, difficulty breathing, congested, cough, tightness of throat
- ⊛ **Stomach:** Discomfort, nausea, vomiting, abdominal cramps, diarrhea
- ⊛ **Skin:** Hives, swelling, rash

When you see any of the above symptoms it is important to initiate the following plan of care:

- * If possible flush the area with large amounts of water.
- * diphenhydramine: Dosage _____ to be given _____
- * epinephrine: Dosage _____ to be given when the following symptoms occur:

NOTE: If epinephrine is given emergency medical services (911) must be accessed immediately.

Provider – please check Level of Assistance:

Independent Student: No assistance is needed from school staff except during emergencies. May self-administer and carry medicine. A back up supply may also be kept in Health Office if desired.

I attest this student has demonstrated to me they can self-administer the medication(s) ordered above safely & effectively, and may carry and use this medication independently at any school/school sponsored activity with no supervision by school staff except in emergencies.

Supervised Student: Adult school staff trained by RN may assist student when taking medication ordered above. The medication will be kept in Health Office or with adult school staff at school/school sponsored activity (i.e. field trips).

Health Care Provider's Signature: _____ Date: _____

Health Care Provider's Name: _____ Phone: _____


Preferred Hospital: _____

Parent / Guardian Signature: _____ Date: _____

The Parent / Guardian signature authorizes the school to share this information with school staff on a "need-to-know" basis. In the event of an emergency care will be initiated and parents will be contacted.

Directions for EpiPen® Use

- Green label for small child.
 - Yellow label for larger child or adult.
1. Hold firmly with orange tip pointing downward.
 2. Remove blue safety release.
 3. Swing and push orange tip firmly into mid-outer thigh (through clothing if necessary) until you hear a 'click'.
 4. Hold in place for 10 seconds.
 5. Massage the injection site for 10 seconds
 6. Once EpiPen® is used, call 911/EMS. Take the used EpiPen® to the emergency room with you.




Directions for Auvi-Q® Use

- Blue for small child.
 - Orange for larger child or adult.
1. Pull Auvi-Q® from the outer case, audio instructions begin.
 2. Pull off red safety guard
 3. Place black end against outer thigh (through clothing if necessary), press firmly and hold for 5 seconds.
 4. The Auvi-Q® makes a distinct click and hiss when activated, this is normal.
 5. Once Auvi-Q® is used, call 911/EMS. Take the used Auvi-Q® to the emergency room with you.



Directions for Adrenaclick®

- Orange label for small child.
 - Yellow label for larger child or adult.
1. Remove the outer case.
 2. Remove grey caps labeled "1" and "2".
 3. Place red rounded tip against mid-outer thigh.
 4. Press down hard until needle penetrates.
 5. Hold for 10 seconds. Remove from thigh.
 6. Once Adrenaclick® is used, call 911/EMS. Take the used Adrenaclick® to the emergency room with you.



Directions for Teva® – Generic EpiPen

- Green label for small child.
 - Yellow label for larger child or adult
1. Twist off green or yellow cap in direction of "twist arrow" to remove cap.
 2. Grasp injector in your hand with orange tip pointing down. With your other hand pull off blue safety release.
 3. Swing and push orange tip against mid-outer thigh till click is heard. Hold in place 3 seconds.
 4. Once EpiPen is used, call 911/EMS. Take used EpiPen to the ER or give to EMS.

